

SOME GENERAL PROPERTIES OF STEIN AG-GROUPOIDS AND STEIN-TEST

M. RASHAD^A, I. AHMAD^{A,*}, AND M. SHAH^B

ABSTRACT. A groupoid S that satisfies the left invertive law, $ab \cdot c = cb \cdot a$ is called an AG-groupoid. We extend this concept to introduce a Stein AG-groupoid. We prove the existence of this type of AG-groupoid by providing some non-associative examples. We also explore some of the basic and general properties of these AG-groupoids and find relations with other known subclasses of AG-groupoids. We present a table of enumeration for these AG-groupoids up to order 6 and further categorize into associative and non-associative. We also develop a method to test an arbitrary AG-groupoid for this new class. Further, we also characterize these AG-groupoids by the properties of their ideals.

1. Introduction and Preliminaries

An AG-groupoid S is a non-associative algebraic structure that generalizes the commutative semigroup in general, and satisfies the left invertive law, $ab \cdot c = cb \cdot a$. One can easily verify that every AG-groupoid satisfies the medial property, $ab.cd = ac.bd$. A groupoid S is called a Stein groupoid if it satisfies the Stein identity $a \cdot bc = bc \cdot a \forall a, b, c \in S$ [2, 5]. We adjoin this property to AG-groupoid to introduce a Stein AG-groupoid. AG-groupoids have been enumerated up to order 6 in [4]. Using the same techniques and data we also enumerate our Stein AG-groupoids up to order 6. Table 1 provides the enumeration of our Stein AG-groupoids. In this note we provide various non-associative examples for this class. It is worth mentioning that various other new classes of AG-groupoids have been recently discovered, enumerated and discussed by various authors [9, 10, 11, 14, 17, 18]. We also define and list various examples of Stein AG-groupoid in Section 2. Section 3 provides a procedure of testing an arbitrary AG-groupoid for a Stein AG-groupoid. In Section 4 we discuss some relations of Stein AG-groupoid with

Key words and phrases. AG-groupoid, Stein-groupoid, Stein AG-test, locally associative AG-groupoid, ideals, nuclear square AG-groupoid.

2010 Mathematics Subject Classification: 20N02, 20M10

*Corresponding author.

already known classes of AG-groupoids[15, 16, 19] and investigate its general properties. While in Section 5 we characterize Stein AG-groupoids by the properties of their ideals.

In the following we list some of the already known classes of AG-groupoids with their identities that will be used frequently in the rest of this article.

Definition 1. [1] *A groupoid S is called an AG-groupoid if $ab \cdot c = cb \cdot a$ for all $a, b, c \in S$.*

Definition 2. [3, 8, 9] *An AG-groupoid S is called —*

- (a) — *a left nuclear square AG-groupoid if $(a^2b)c = a^2(bc)$, $\forall a, b, c \in S$.*
- (b) — *a middle nuclear square AG-groupoid if $(ab^2)c = a(b^2c)$, $\forall a, b, c \in S$.*
- (c) — *a right nuclear square AG-groupoid if $(ab)c^2 = a(bc^2)$, $\forall a, b, c \in S$.*
- (d) — *a right alternative AG-groupoid if $a(bb) = (ab)b$, $\forall a, b \in S$.*
- (e) — *a Bol*-AG-groupoid if $a(bc \cdot d) = (ab \cdot c)d$, $\forall a, b, c, d \in S$.*
- (f) — *a paramedial AG-groupoid if $ab \cdot cd = db \cdot ca$, $\forall a, b, c, d \in S$.*

Definition 3. [6, 8, 9, 13] *An AG-groupoid S is called —*

- (a) — *an AG*-groupoid if $ab \cdot c = b \cdot ac$, $\forall a, b, c \in S$.*
- (b) — *a T_f^4 -AG-groupoid if $ab = cd \Rightarrow ad = cb$, $\forall a, b, c, d \in S$.*
- (c) — *a T_b^4 -AG-groupoid if $ab = cd \Rightarrow da = bc$, $\forall a, b, c, d \in S$.*
- (d) — *an AG-3-band if $a(aa) = a$, $\forall a \in S$.*
- (e) — *a cancellative AG-groupoid if $ax = bx$, and $xa = ya \Rightarrow a = b$, $\forall a, b \in S$.*
- (f) — *a medial AG-groupoid if $ab \cdot cd = ac \cdot bd$, $\forall a, b, c, d \in S$.*
- (g) — *locally associative if $a(aa) = (aa)a$, $\forall a \in S$.*

Order	3	4	5	6
Non-associative AG-groupoids	8	269	31467	40104513
Associative AG-groupoids	12	62	446	7510
Non-associative Stein AG-groupoids	0	0	16	931
Associative Stein AG-groupoids	5	14	46	173

TABLE 1. Enumeration of Stein AG-groupoids up to order 6.

2. Stein AG-groupoid

In this section we define a Stein AG-groupoid and give a non-associative example of lowest order to show its existence.

Definition 4. An AG-groupoid S is called a Stein AG-groupoid if,

$$(2.1) \quad a(bc) = (bc)a \quad \forall a, b, c \in S$$

Example 1. Let $S = \{1, 2, 3, 4, 5\}$. Then it is easy to verify that S is a Stein AG-groupoid.

$\cdot$	1	2	3	4	5
1	1	1	1	1	1
2	1	1	1	1	1
3	1	1	1	2	2
4	1	1	2	2	3
5	1	1	2	2	3

3. Stein AG-test

In this section we discuss a procedure by [12] to check an arbitrary AG-groupoid $(G, \cdot)$ for a Stein-AG-groupoid. To this end we define the following binary operations.

$$(3.1) \quad a \circ b = a \cdot bx$$

$$(3.2) \quad a \triangle b = bx \cdot a$$

The identity $a \cdot bx = bx \cdot a$ holds if,

$$(3.3) \quad a \circ b = a \triangle b$$

To construct table of the operation “ $\circ$ ” for any fixed $x \in G$, we multiply turn by turn elements of index column of the “ $\cdot$ ” table from left with the elements of x-column of the “ $\cdot$ ” table. Similarly the table of operation “ $\triangle$ ” for any fixed $x \in G$ is obtained by presenting x-column of the “ $\cdot$ ” table as index column and multiplying it with elements of index row of the “ $\cdot$ ” table from the left. If the tables for the operation “ $\circ$ ” and “ $\triangle$ ” coincides for all $x \in G$, then 2.1 holds and the AG-groupoid is a Stein AG-groupoid in this case. We illustrate this procedure in the following example.

Example 2. Verify the following AG-groupoid G for a Stein AG-groupoid.

$\cdot$	1	2	3	4	5
1	1	1	1	1	1
2	1	1	1	1	1
3	1	1	1	1	1
4	1	1	1	1	2
5	1	1	4	2	1

[illegible]

TABLE 2.

Extend the above table in a way as described above. The tables to the right of original “ $\cdot$ ” table are constructed for the operation “ $\circ$ ” and the downwards tables are constructed for the operation “ Δ ”. It is clear from the extended table that the downward tables and the tables on the right coincide so G is a Stein AG-groupoid.

4. Characterization of Stein AG-groupoids

In this section we find the relations of Stein AG-groupoids with already other known classes of AG-groupoids. We start with the following:

Lemma 1. *[3]AG^{**}-groupoid is Bol^{*}-AG-groupoid.*

Lemma 2. *[3]Bol^{*}-AG-groupoid is paramedial AG-groupoid.*

Lemma 3. *[3]AG^{**}-groupoid is left nuclear square AG-groupoid.*

Proposition 1. *Let S be a Stein AG-groupoid, then*

- (a) *S is locally associative AG-groupoid.*
- (b) *S is right alternative AG-groupoid.*
- (c) *S is AG^{**}-groupoid.*
- (d) *S is Bol^{*}-AG-groupoid.*
- (e) *S is paramedial AG-groupoid.*
- (f) *S is left nuclear square AG-groupoid.*
- (g) *S is right nuclear square AG-groupoid.*
- (h) *S is middle nuclear square AG-groupoid.*
- (i) *S is nuclear square AG-groupoid.*

Proof. Let S be a Stein AG-groupoid, and $a, b, c \in S$.

- (a) Since S is a Stein AG-groupoid, thus $a(bc) = (bc)a$, $\forall a, b, c \in S$. Thus by replacing b and c by a , we get $a(aa) = (aa)a$. Hence S is locally associative AG-groupoid.
- (b) Since $\forall a, b \in S$ we have, $ab.b = (bb)a = a(bb) \Rightarrow (ab)b = a(bb)$. Hence S is right alternative AG-groupoid.
- (c) Since S is Stein, thus $\forall a, b, c \in S$, $a(bc) = (bc)a = (ac)b = b(ac) \Rightarrow a(bc) = b(ac)$. Hence S is AG^{**}-groupoid.
- (d) The result follows from Lemma 1.
- (e) Follows from Lemma 2.
- (f) Follows from Lemma 3.
- (g) We show that S is right nuclear square AG-groupoid. Let $a, b, c \in S$. Then

$$\begin{aligned} (ab)c^2 &= (cc \cdot b)a = (b \cdot cc)a = (bc^2)a = \\ &= a(bc^2) \Rightarrow (ab)c^2 = a(bc^2). \end{aligned}$$

Hence the result is proved.

(h) Let $a, b, c \in S$. Then

$$\begin{aligned} a \cdot b^2 c &= (bb.c)a = (c \cdot bb)a = (cb^2)a = \\ &= ab^2 \cdot c \Rightarrow a \cdot b^2 c = ab^2 \cdot c. \end{aligned}$$

Hence S is middle nuclear square AG-groupoid.

(i) Follows by (f), (g) and (h). ■

Theorem 1. *A Stein AG-groupoid S is a semigroup if any of the following hold.*

- (i) S is AG^* -groupoid.
- (ii) S is AG-3-band.
- (iii) S is T^4 -AG-groupoid.
- (iv) S has a cancellative element.

Proof. Let S be a Stein AG-groupoid.

(i) Assume that S is AG^* -groupoid, and $a, b, c \in S$. Then using Definitions 1, 3 and 4, we have

$$ab \cdot c = b.ac = ac.b = bc \cdot a = a \cdot bc \Rightarrow ab \cdot c = a \cdot bc.$$

(ii) Let S be an AG-3-band, and $a, b, c \in S$. Then by Definitions 1, 3 and 4, we get

$$\begin{aligned} ab \cdot c &= cb \cdot a = (cb)(aa \cdot a) = (c \cdot aa)(ba) = \\ &= (ba)(c \cdot aa) = (bc)(a \cdot aa) = (bc)(aa \cdot a) = \\ &= bc \cdot a = a \cdot bc \Rightarrow ab \cdot c = a \cdot bc. \end{aligned}$$

(iii) Let S be a T^4 -AG-groupoid, and $a, b, c \in S$. Then by Definitions 1, 3 and 4, we have

$$\begin{aligned} ab \cdot c &= cb.a = a.cb \Rightarrow ab.cb = ac \Rightarrow ac.bb = ac \\ &\Rightarrow ac \cdot c = a \cdot bb = bb \cdot a = ab \cdot b \Rightarrow ac \cdot b = ab \cdot c \\ &\Rightarrow bc \cdot a = ab \cdot c \Rightarrow a \cdot bc = ab \cdot c. \end{aligned}$$

(iv) Let S has a left cancellative element x , and $a, b, c \in S$. Then by Definitions 1, 3 and 4, it is clear that

$$\begin{aligned} x(a \cdot bc) &= (a.bc)x = (x.bc)a = (bc.x)a = (xc \cdot b)a = \\ &= ab \cdot xc = xc \cdot ab = (ab \cdot c)x = x(ab \cdot c) \\ &\Rightarrow x(a \cdot bc) = x(ab \cdot c) \Rightarrow a \cdot bc = ab \cdot c. \end{aligned}$$

- (v) Let S has a right cancellative element x , and $a, b, c \in S$. Then by Definitions 1, 3 and 4, we have

$$\begin{aligned} (a \cdot bc)x &= (x.bc)a = (bc.x)a = ax \cdot bc = ab \cdot xc = xc \cdot ab = \\ &= (ab \cdot c)x \Rightarrow (a \cdot bc)x = (ab \cdot c)x \Rightarrow a \cdot bc = ab \cdot c. \end{aligned}$$

Hence S is a semigroup. ■

5. Ideals in Stein AG-groupoids

A subset A of the AG-groupoid S is a left (resp. right) ideal of S if,

$$(5.1) \quad SA \subseteq A \text{ (resp. } AS \subseteq A)$$

A is a two sided ideal or simply an ideal of S if it is both left and right ideal of S .

Let S be an AG-groupoid and $A, B \subseteq S$, then A and B are called left connected if $SA \subseteq B$ and $SB \subseteq A$. Similarly, if S is an AG-groupoid and $A, B \subseteq S$, then A and B are called right connected sets if $AS \subseteq B$ and $BS \subseteq A$ and are called connected if are both left and right connected.

Lemma 4. *If S is a Stein AG-groupoid and $A, B \subseteq S$. Then AB and BA are right and left connected sets.*

Proof. Let S be a Stein AG-groupoid. Then by Definitions 1 and 4, we get

$$S(AB) = (AB)S = (SB)A \subseteq BA \Rightarrow S(AB) \subseteq BA.$$

Similarly,

$$S(BA) = (BA)S = SA \cdot B \subseteq AB \Rightarrow S(BA) \subseteq AB.$$

Hence AB and BA are left connected sets. Now using Definition 1, we have

$$(AB)S = (SB)A \subseteq BA \Rightarrow (AB) \subseteq BA.$$

Similarly,

$$(BA)S = SA \cdot B \subseteq AB \Rightarrow S(BA) \subseteq AB.$$

Hence AB and BA are right connected sets. ■

Proposition 2. *If S is a Stein AG-groupoid and L is left ideal of S then L^2 is an ideal of S .*

Proof. Since $S(L^2) = (LL)S = (SL)L \subseteq L \cdot L = L^2$ and $(L^2)S = SL \cdot L \subseteq LL = L^2$. Hence L^2 is an ideal. ■

Theorem 2. *If S is a Stein AG-groupoid. Then for any fixed a in S*

- (a) aS is an ideal of S .
- (b) $a(Sa)$ is minimal ideal of S .
- (c) $(aS)a$ is an ideal of S .

Proof. Let S be a Stein AG-groupoid. Then

- (a) Let a be any fixed element of S then,

$$\begin{aligned} S(aS) &= \bigcup_{x,y \in S} x(ay) = \bigcup_{x,y \in S} (ay)x = \\ &= \bigcup_{x,y \in S} xy \cdot a = \bigcup_{x,y \in S} a \cdot xy \subseteq aS \\ \Rightarrow S(aS) &\subseteq aS. \end{aligned}$$

Thus aS is a left ideal of S . Similarly,

$$\begin{aligned} (aS)S &= \bigcup_{x,y \in S} (ax)y = \bigcup_{x,y \in S} yx \cdot a = \bigcup_{x,y \in S} a \cdot yx \subseteq aS \\ \Rightarrow (aS)S &\subseteq aS. \end{aligned}$$

Thus aS is a right ideal of S . Equivalently aS is an ideal.

- (b) Again by Definitions 1, 4 and Proposition 1(c), we get

$$\begin{aligned} S(a(Sa)) &= \bigcup_{x,y \in S} x(a \cdot ya) = \bigcup_{x,y \in S} x(ya \cdot a) = \bigcup_{x,y \in S} x(aa \cdot y) = \\ &= \bigcup_{x,y \in S} (aa \cdot y)x = \bigcup_{x,y \in S} xy \cdot aa = \\ &= \bigcup_{x,y \in S} a(xy \cdot a) \subseteq a(Sa) \Rightarrow S(a(Sa)) \subseteq (a(Sa)). \end{aligned}$$

Thus $a(Sa)$ is a left ideal of S . Similarly by Definition 1, 4 and Proposition 1(c), we have

$$\begin{aligned} (a(Sa))S &= \bigcup_{x,y \in S} (a \cdot xa)y = \bigcup_{x,y \in S} (xa \cdot a)y = \\ &= \bigcup_{x,y \in S} (aa \cdot x)y = \bigcup_{x,y \in S} yx \cdot aa = \\ &= \bigcup_{x,y \in S} a(yx \cdot a) \subseteq a(Sa) \Rightarrow S(a(Sa)) \subseteq (a(Sa)) \end{aligned}$$

Hence $a(Sa)$ right ideal of S . Thus $a(Sa)$ is an ideal of S . Let L be an ideal of S , such that $a \in L$. Then

$$a(Sa) \subseteq L(SL) \subseteq LL \subseteq L \text{ also } a(Sa) \subseteq L(SL) = (SL)L \subseteq LL \cdot S \subseteq LS \subseteq L.$$

Hence $a(Sa)$ is minimal ideal of S .

(c) Now using Definitions 1 and 4, we have

$$\begin{aligned}
 S((aS)a) &= \bigcup_{x,y \in S} x(ay \cdot a) = \bigcup_{x,y \in S} x(a \cdot ay) = \bigcup_{x,y \in S} (a \cdot ay)x = \\
 &= \bigcup_{x,y \in S} (x \cdot ay)x = \bigcup_{x,y \in S} (ay \cdot x)a = \bigcup_{x,y \in S} (xy \cdot a)a = \\
 &\Rightarrow S((aS)a) = \bigcup_{x,y \in S} (a \cdot xy)a \subseteq (aS)a \subseteq (aS)a.
 \end{aligned}$$

So $(aS)a$ is left ideal of S . Similarly by Definition 4, we have

$$\begin{aligned}
 ((aS)a)S &= \bigcup_{x,y \in S} (ax \cdot a)y = \bigcup_{x,y \in S} (a \cdot ax)y = \bigcup_{x,y \in S} (y \cdot ax)a = \\
 &= \bigcup_{x,y \in S} (ax \cdot y)a = \bigcup_{x,y \in S} (yx \cdot a)a = \\
 &= \bigcup_{x,y \in S} (a \cdot yx)a \subseteq (aS)a \Rightarrow S((aS)a) \subseteq (aS)a.
 \end{aligned}$$

Hence $a(Sa)$ is right ideal. Thus $(aS)a$ is an ideal of S . ■

Theorem 3. *Let L be a left ideal and R be a right ideal of a Stein-AG-groupoid S , such that $x = x^2$ and $y = y^2$ for some $x, y \in S$. Then*

- (i) $xL = L \cap xS$
- (ii) $Rx = Sx \cap R$
- (iii) $Sy \cap xS = x(Sy)$

Proof. Let L be a left ideal and R be a right ideal of S , and $x, y \in S$ such that $x = x^2$ and $y = y^2$. Then

- (i) Let $t \in xL$, so $t = xl$ for some $l \in L$. Since $l \in L$ and $x \in S$, then $xl \in L$, so that $t \in L$. But $l \in L \subseteq S$, therefore $l \in S$, and so $xl = t \in xS$. Hence $t \in L \cap xS$, that is

$$(5.2) \quad xL \subseteq L \cap xS$$

Conversely, let $u \in L \cap xS$, then $u \in L$ and $u \in xS$, so $u = xs$ for some $s \in S$. Now by Definition 4, Proposition 1 (c) and the assumption, we get

$$\begin{aligned}
 u &= xs = (xx)s = s(xx) = x(sx) = \\
 &= x(s \cdot xx) = x(xx \cdot s) = x(xs) = \\
 &= xu \in xL \Rightarrow u \in xL.
 \end{aligned}$$

Therefore,

$$(5.3) \quad L \cap xS \subseteq xL$$

Hence by (5.2) and (5.3), we have $L \cap xS = xL$.

- (ii) Let $b \in Rx$, so $b = rx$ for some $r \in R$. Since $r \in R$ and $x \in S$, then $rx \in R$, so that $b \in R$. But $r \in R \subseteq S$, then $r \in S \Rightarrow rx = b \in Sx$, hence $b \in R \cap Sx$, this implies that

$$(5.4) \quad Rx \subseteq R \cap Sx$$

Conversely, let $t \in Sx \cap R$, then $t \in Sx$ and $t \in R$. Since $t \in Sx$, so $t = sx$ for some $s \in S$. Now by assumption and Definition 1, 4, we have

$$t = sx = s(xx) = (xx)s = (sx)x = tx \in Rx \Rightarrow t \in Rx.$$

Therefore,

$$(5.5) \quad Sx \cap R \subseteq Rx$$

Hence by equations 5.4 and 5.5, we have $Sx \cap R = Rx$.

- (iii) If $a \in Sy \cap xS$, then for some $s, s' \in S$, we have $a = sy$ and $a = xs'$. Now by Definitions 1, 4 and assumption, we have $ay = sy \cdot y = yy \cdot s = s \cdot yy = sy = a$. and

$$\begin{aligned} a &= xs' = s'x \cdot x = s'x \cdot xx = \\ &= x(s'x \cdot x) = x(xx \cdot s') = x \cdot xs' = xa. \end{aligned}$$

Therefore $ay = xa = a$, and so again by Definitions 1, 4 and assumption, we have

$$\begin{aligned} x(ay) &= x(xa) = xx \cdot xa = xa \cdot xx = xx \cdot ax = \\ &= x \cdot ax = ax \cdot x = xx \cdot a = xa = a. \end{aligned}$$

Thus $a = x(ay) \Rightarrow a \in x(Sy)$. Therefore,

$$(5.6) \quad Sy \cap xS \subseteq x(Sy)$$

Conversely, if $a \in x(Sy)$, then by Definitions 1, 4 and assumption, we have

$$\begin{aligned} a &= x(sy) = x(s \cdot yy) = x(yy \cdot s) = \\ &= x(sy \cdot y) = x(y \cdot sy) = (y \cdot sy)x = (x \cdot sy)y = ay, \end{aligned}$$

and

$$a = x(sy) = xx \cdot sy = (sy \cdot x)x = (x \cdot sy)x = x(x \cdot sy) = xa.$$

That is $a \in Sy \cap xS$. Therefore,

$$(5.7) \quad x(Sy) \subseteq Sy \cap xS$$

Hence by 5.6 and 5.7, we have $x(Sy) = Sy \cap xS$. ■

REFERENCES

- [1] M.A. Kazim and M. Naseerudin, On almost semigroups, Portugaliae Mathematica, 36(1) (1977).
- [2] M.J. Pelling and D.G. Rogers, Some (2,n) combinatorial Stein groupoids, aequationes mathematicae, Volume 21, Issue 1, pp 225-235 (1980).
- [3] M. Shah, A theoretical and computational investigation of AG-groups, PhD thesis, Quaid-i-Azam University Islamabad, (2012).
- [4] A. Distler, M. Shah and V. Sorge, Enumeration of AG-groupoids, Lecture Notes in Computer Science, Volume 6824/2011, 1-14 (2011).
- [5] J. Denes and A.D. Keedwell, Latin Squares and their applications, Academic Press New York and London, 1974.
- [6] Q. Mushtaq and S.M. Yusuf, On Locally Associative LA-Semigroup, J. Nat. Sci. Math., XIX(1), 57-62 (1979).
- [8] J. R. Cho, J. Jezek and T. Kepka, Paramedial Groupoids, Czechoslovak Mathematical Journal, 49 (124) 1996.
- [9] M. Shah, I. Ahmad, and A. Ali, On introduction of new classes of AG-groupoids, Res. J. Recent Sci., Vol. 2(1), 2013, 67-70.
- [10] M. Shah, I. Ahmad, and A. Ali, Discovery of new classes of AG-groupoids, Res. J. Recent Sci., Vol. 1(11), 2012, 47-49.
- [11] M. Rashad, I. Ahmad, Amanullah and M. Shah, On relations between right alternative and nuclear square AG-groupoids, Int. Mathematica Forum, Vol. 8(5), 237-243, 2013.
- [12] P.V. Protic and N. Stevanovic, AG-test and some general properties of Abel-Grassmann's groupoids, PU. M. A, 6 (1995), 371 - 383.
- [13] I. Ahmad, M. Rashad and M. Shah, Some new result on T^1 , T^2 and T^4 -AG-groupoids, Research Journal of Recent Sciences, Vol. 2(3), 64-66, (2013).
- [14] M. Khan, Faisal, V. Amjad, On some classes of Abel-Grassmann's groupoids. J. Adv. Res. Pure Math. 3(4), 109-119 (2011)
- [15] M. Rashad, I. Ahmad and M. Shah, Left transitive AG-groupoids, [www.arXiv.org 1402.5296](http://www.arXiv.org/abs/1402.5296).
- [16] M. Rashad, I. Ahmad, and M. Shah, Bi-commutative AG-groupoids, [www.arXiv.org. 1403.2564](http://www.arXiv.org/abs/1403.2564)
- [17] Amanullah, M. Rashad, I. Ahmad, M. Shah, On Modulo AG-groupoids, [arXiv: 1403.2304](http://www.arXiv.org/abs/1403.2304)
- [18] Amanullah, M. Rashad, I. Ahmad, M. Shah, M. Yousaf, Modulo Matrix AG-groupoids and Modulo AG-groups [arXiv:1403.2304](http://www.arXiv.org/abs/1403.2304)
- [19] P. Yiarayong, On Semiprime and Quasi Semiprime ideals in AG-groupoids, KKU Res. J. 2013; 18(6): 893-896

E-mail address: rashad@uom.edu.pk

E-mail address: iahmaad@hotmail.com

A. DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MALAKAND, CHAKDARA DIR(L), PAKISTAN.

E-mail address: shahmaths_problem@hotmail.com

B. DEPARTMENT OF MATHEMATICS, GOVERNMENT POST GRADUATE COLLEGE
MARDAN, PAKISTAN.